# **National Parks and Wildlife Service**

**Conservation Objectives Series** 

## Knockalongy and Knockachree Cliffs SAC 001669



An Roinn Tithíochta, Rialtais Áitiúil agus Oidhreachta Department of Housing, Local Government and Heritage National Parks and Wildlife Service, Department of Housing, Local Government and Heritage,

90 King Street North, Dublin 7, D07 N7CV, Ireland.

Web: www.npws.ie E-mail: nature.conservation@chg.gov.ie

Citation:

NPWS (2020) Conservation Objectives: Knockalongy and Knockachree Cliffs SAC 001669. Version 1. National Parks and Wildlife Service, Department of Housing, Local Government and Heritage.

Series Editor: Rebecca Jeffrey ISSN 2009-4086

#### Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance
- exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

• population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and

• the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and

• there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

#### Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.

2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.

3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.

4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.

5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

### Qualifying Interests

\* indicates a priority habitat under the Habitats Directive

001669 Knockalongy and Knockachree Cliffs SAC

6985 Killarney Fern Vandenboschia speciosa

### Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

### **NPWS Documents**

Year :	2015		
Title :	Monitoring methods for the Killarney Fern (Trichomanes speciosum Willd.) in Ireland		
Author :	Ní Dhúill, E.; Smyth, N.; Waldren, S.; Lynn, D.		
Series :	Irish Wildlife Manuals, No. 82		
Year :	2019		
Title :	The Status of EU Protected Habitats and Species in Ireland. Volume 3: Species Assessments		
Author :	NPWS		
Series :	Conservation assessments		
Year :	in prep.		
Title :	Monitoring and assessment of Killarney Fern ( <i>Vandenboschia speciosa</i> (Willd.) Kunkel) in Ireland, 2015-2018		
Author :	Ní Dhúill, E.; O'Neill, F.H.; Hodd, R.		
Series :	Irish Wildlife Manuals		

### Conservation Objectives for : Knockalongy and Knockachree Cliffs SAC [001669]

### 6985 Killarney Fern Vandenboschia speciosa

To maintain the favourable conservation condition of Killarney Fern in Knockalongy and Knockachree Cliffs SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Occurrence	No loss in geographical spread of populations, subject to natural processes	The population of Killarney fern ( <i>Vandenboschia</i> <i>speciosa</i> [formerly <i>Trichomanes speciosum</i> ; species code 1421]) is currently known from locations in Knockalongy and Knockachree Cliffs SAC within hectad G52. Exact locations are not mapped here or account of the threat posed by illegal collecting. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Number of populations	Number	No decline, subject to natural processes	One population of the species has been recorded in the SAC since 1960. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Number of colonies	Number	No decline, subject to natural processes	Two colonies of the species have been recorded in the population in the SAC since 1960. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Population: life- cycle stage	Type (sporophyte or gametophyte)	Maintain life-cycle stage composition of the population, subject to natural processes	One of the two colonies recorded since 1960 is composed of sporophytes (frond stage) with coexisting gametophytes (filamentous stage) and one is composed of gametophytes only. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Population size: area of occupancy	Square metres	No decline, subject to natural processes	Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Population size: living sporophyte fronds	Number	No decline, subject to natural processes	Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Population structure: young and unfurling fronds	Occurrence	Young (not fully expanded) and/or unfurling (crozier) fronds present in populations previously observed to have these, subject to natural processes	Young and/or unfurling fronds have been recorded from Knockalongy and Knockachree Cliffs SAC. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Population structure: fertile fronds	Occurrence	Fertile fronds present in populations previously observed to have these, subject to natural processes	Fertile fronds have not been recorded from the SAC. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Population structure: juvenile sporophyte fronds emerging from gametophytes	Number	No decline, subject to natural processes	Juvenile sporophyte fronds emerging from gametophytes have not been recorded from the SAC. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Habitat extent	Hectares	No loss of suitable habitat, subject to natural processes	The species grows in deeply shaded, humid situations - dripping caves, overhangs and crevices on cliffs, rocky slopes, by waterfalls, in stream ravines and gullies, on rock or soil banks in woodlands and, occasionally, under fallen trees and on the floor of damp woodlands. Whilst also occurring in these habitats, the gametophyte stage can grow in drier areas that do not suit the sporophyte. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files

Hydrological conditions: wet/damp microhabitats	Occurrence	Maintain hydrological conditions at the locations of the known populations- visible water source, with dripping or seeping water present and/or substrate wet/damp to touch, subject to natural processes	Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Hydrological conditions: relative humidity	Percentage	Maintain relative humidity levels at known colonies at not less than 80%, subject to natural processes	Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Hydrological conditions: desiccated fronds	Number	No increase, subject to natural processes	Presence of desiccated sporophyte fronds and gametophyte mats is indicative of unsuitable conditions. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Light levels: shading	Shade index score	colonies; at least 5 for open upland sporophyte-	Shade Index: 4. Moderate shade, e.g. light-medium deciduous canopy with sun flecks. 5. Permanently shaded from direct sunlight but otherwise open to sky. 6. Deep woodland (e.g. coniferous or in ravine) shade, no sun flecks. 7. Perpetual deep shade, e.g. cave entrance, beneath boulder. Woodland colonies have not been recorded from the SAC to date. The species occurs in permanent shade in Knockalongy and Knockachree Cliffs SAC. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files
Woodland canopy cover	Percentage	No loss of woodland canopy at, or in the vicinity of, the locations of known populations and canopy cover here maintained at more than 33%, subject to natural processes	of its habitat requirements, particularly with regard to maintenance of sufficient canopy cover. Woodland colonies have not been recorded from the
Invasive species	Occurrence	Maintain absence of invasive non-native and vigorous native plant species at the locations of the known population or, if present, maintain vegetation cover of these at less than 10%, taking into account the habitat requirements of <i>V.</i> <i>speciosa</i>	In order to avoid negative impacts on the Killarney fern ( <i>Vandenboschia speciosa</i> ), its habitat requirements (site hydrology, relative humidity, canopy cover, shading levels, etc.) must be taken into account in locations that are subject to or proposed for management actions to control invasive non-native and/or vigorous native plant species. Based on Ní Dhúill et al. (2015, in prep.), NPWS (2019) and NPWS internal files

